

Please send by the announced due date to Glen Cowan, Physics Dept., Royal Holloway, University of London, Egham, Surrey, TW20 0EX, or e-mail to [g.cowan@rhul.ac.uk](mailto:g.cowan@rhul.ac.uk).

**Exercise 2.1:** Suppose the independent random variables  $x_1$  and  $x_2$  have means  $\mu_1 = \mu_2 = 10$  and variances  $\sigma_1^2 = \sigma_2^2 = 1$ . Use error propagation to find the variance of

$$y = \frac{x_1^2}{x_2}. \quad (1)$$

Comment on the validity of the procedure if one had  $\mu_2 = 1$ .

**Exercise 2.2:** Suppose the random variable  $x$  is uniformly distributed in the interval  $[\alpha, \beta]$ , with  $\alpha, \beta > 0$ . Find the expectation value of  $1/x$ , and compare the answer to  $1/E[x]$  using  $\alpha = 1, \beta = 2$ .

**Exercise 2.3:** Suppose two independent random variables  $x$  and  $y$  are both uniformly distributed between zero and one, i.e. the p.d.f.  $g(x)$  is given by

$$g(x) = \begin{cases} 1 & 0 < x < 1 \\ 0 & \text{otherwise,} \end{cases} \quad (2)$$

and similarly for the p.d.f.  $h(y)$ .

(a) Show that the p.d.f.  $f(z)$  for  $z = xy$  is

$$f(z) = \begin{cases} -\log z & 0 < z < 1 \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

by defining an additional function,  $u = x$ . First, find the joint p.d.f. of  $z$  and  $u$ . Integrate this over  $u$  to find the p.d.f. for  $z$ .

(b) Show that the cumulative distribution of  $z$  is

$$F(z) = z(1 - \log z). \quad (4)$$

**Exercise 2.4:** Consider the exponential p.d.f.,

$$f(x; \xi) = \frac{1}{\xi} e^{-x/\xi}, \quad x \geq 0. \quad (5)$$

(a) Show that the corresponding cumulative distribution is given by

$$F(x) = 1 - e^{-x/\xi}, \quad x \geq 0. \quad (6)$$

(b) Show that the conditional probability to find a value  $x$  between  $x_0$  and  $x_0 + x'$  given that  $x > x_0$  is equal to the (unconditional) probability to find  $x$  less than  $x'$ , i.e.

$$P(x \leq x_0 + x' | x \geq x_0) = P(x \leq x'). \quad (7)$$

(c) Cosmic ray muons produced in the upper atmosphere enter a detector at sea level, and some of them come to rest in the detector and decay. The time difference  $t$  between entry into the detector and decay follows an exponential distribution, and the mean value of  $t$  is the mean lifetime of the muon (approximately  $2.2 \mu\text{S}$ ). Explain why the time that the muon lived prior to entering the detector does not play a role in determining the mean lifetime.